

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.

Iatrogenic pulmonary embolism with cyanoacrylate: to remove or to leave?

Authors: Arkadiusz Pietrasik, Aleksandra Gąsecka, Dominika Chojecka, Jakub Pytlos, Bartosz Rymuza, Renata Głowczyńska, Marta Banaszkiewicz, Szymon Darocha, Marcin Kurzyna

Article type: Clinical vignette

Received: February 22, 2021.

Accepted: April 11, 2021.

Published online: April 16, 2021.

ISSN: 0022-9032

e-ISSN: 1897-4279

This is an Open Access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives 4.0 International License ([CC BY-NC-ND 4.0](https://creativecommons.org/licenses/by-nc-nd/4.0/)), allowing third parties to download articles and share them with others, provided the original work is properly cited, not changed in any way, distributed under the same license, and used for noncommercial purposes only. For commercial use, please contact the journal office at kardiologiapolska@ptkardio.pl.

Iatrogenic pulmonary embolism with cyanoacrylate: to remove or to leave?

Short title: Pulmonary embolism with cyanoacrylate: to remove or to leave?

Arkadiusz Pietrasik¹, Aleksandra Gąsecka¹, Dominika Chojecka¹, Jakub Pytlos¹, Bartosz Rymuza¹, Renata Głównczyńska¹, Marta Banaszekiewicz², Szymon Darocha², Marcin Kurzyna²

¹ 1st Chair and Department of Cardiology, Medical University of Warsaw, Warsaw, Poland

² Department of Pulmonary Circulation, Thromboembolic Diseases and Cardiology, Centre of Postgraduate Medical Education, European Health Centre Otwock, Poland

Corresponding author:

Aleksandra Gąsecka, MD, PhD

1st Chair and Department of Cardiology, Medical University of Warsaw

Banacha 1a, 02-097, Warsaw, Poland

E-mail: aleksandra.gasecka@wum.edu.pl

Telephone: +48 22 599 19 51

Conflicts of Interest: All authors declare that they have no conflicts of interest to disclose.

Word count: 498

A 46-year-old female with autoimmune hepatitis and liver cirrhosis presented with the symptoms of upper gastrointestinal bleeding. Gastroduodenoscopy revealed active hemorrhage from a huge duodenal varix and an endoscopic injection sclerotherapy with cyanoacrylate was performed. Since the bleeding continued, the patient underwent a successful surgical ligation of the bleeding varix the following day.

Twenty days later the patient was re-admitted due to dyspnea, cough and pleuritic chest pain. On physical examination she was tachypneic and hypoxic. Computed tomography angiography (CTA) revealed the presence of disseminated, hyperdense deposits in segmental and subsegmental branches of both pulmonary arteries (**Figure 1A, Video S1**), confirming the previously suspected diagnosis of pulmonary embolism (PE) with cyanoacrylate. A perfusion single-photon-emission computed tomography revealed bilateral wedge-shaped perfusion defects matching several bronchopulmonary segments of both lungs (**Figure 1B, Video S2**). Echocardiography showed no embolic material in heart chambers and no features of right ventricle overload (TAPSE 22 mm), yet pericardial effusion (**Figure 1C, 1D**). The cardiac troponin I and natriuretic peptide concentrations were within normal range.

In contact with blood, cyanoacrylate undergoes rapid polymerization. Large varix size and injected volume increase the risk of embolization via the varix efferent vein into inferior vena cava, right heart chambers and pulmonary arteries. The nature of PE with cyanoacrylate rules out any form of pharmacological therapy, including anticoagulation. The range of surgical treatment options is wide, but choosing an optimal therapy for an individual patient is truly challenging [1]. Hitherto, no consensus regarding the best way of clinical management of iatrogenic cyanoacrylate emboli has been established [2]. For our patient, the accurate management of the PE episode was crucial, since it could impact her overall condition and potentially disqualify her from the liver transplantation. To facilitate immediate decision

making by experts, she was consulted by the local Pulmonary Embolism Response Team (PERT) [3].

Given the hemodynamic stability, our patient was at low-risk of death. The danger of further thromboembolic and septic complications was also identified as unlikely. Consequently, PERT members assessed the risk of interventional therapy to be higher than the risk of death. A decision was made to continue with the conservative therapy, which was followed by a control CTA a month after the episode and a series of endoscopies with varices ligation every 2 months, without further complications. A favorable outcome during the 12-month follow-up period confirmed that the right path of management had been taken.

This report highlights the challenges in the management of iatrogenic PE. Recently, an interesting case of acute PE and right atrial thrombus was presented, which was due to central venous access chemotherapy port migration and required surgical excision [4]. Our patient did not undergo interventional treatment, but only a 12-month follow-up assured us of the patient's recuperation. Hence, in every patient the treatment should be based on the individualized risk stratification to determine whether interventional or conservative approach is more beneficial. Consultation in a multidisciplinary team is an important part of a decision-making process to ensure the optimal clinical management [5].

References:

1. Konstantinides SV, Meyer G, Becattini C, et al. 2019 ESC Guidelines for the diagnosis and management of acute pulmonary embolism developed in collaboration with the European Respiratory Society (ERS). *Eur Respir J*. 2019; 54: 1901647.
2. Asah D, Raju S, Ghosh S, et al. Nonthrombotic Pulmonary Embolism From Inorganic Particulate Matter and Foreign Bodies. *Chest*. 2018; 153: 1249–1265.
3. Rosovsky R, Zhao K, Sista A, et al. Pulmonary embolism response teams: Purpose, evidence for efficacy, and future research directions. *Res Pract Thromb Haemost*. 2019; 3: 315–330.
4. Araszkievicz A, Kurzyna M, Kopeć G, et al. Expert opinion on the creating and operating of the regional Pulmonary Embolism Response Teams (PERT). Polish PERT Initiative. *Cardiol J*. 2019; 26: 623-632.
5. Borowiec A, Kurnicka K, Zieliński D, et al. Acute pulmonary embolism and right atrial thrombus as a complication of the central venous access port device for the delivery of chemotherapy. *Polish Heart Journal*. 2020; 78: 778-779.




Figure 1.

A – Coronal maximum intensity projection showing disseminated, hyperdense cyanoacrylate deposits within the segmental and subsegmental branches of pulmonary arteries to the middle and lower lobe in the right lung and to the upper and lower lobe in the left lung (yellow arrows); there are no residual deposits visible within the right heart;

B – Ventilation (V) and perfusion (Q) single-photon-emission computed tomography slices with multiple mismatched defects in the right and left lung (red arrows);

C, D – Echocardiography showing no embolic material in the heart chambers and no features of right ventricular pressure overload.